


Attorney's Docket: 2002DE446
 Serial No.: 13735490
 Art Unit: 1714
 Response to Office Action, Dated 11/15/2007


This listing of claims will replace all prior versions, and listings, of claims in the application:

1. (Currently Amended) A fuel oil comprising a proportion of mineral oil middle distillate, said fuel oil having a sulfur content of at most 350 ppm, a total aromatics content of at most 22% by weight of said middle distillate, and a proportion of at least one copolymer of ethylene and vinyl esters, said copolymer comprising comonomers

a) bivalent structural units derived from ethylene of the formula 1




b) from 5 to 12 mol% of bivalent structural units of the formula 2



where R¹ is a saturated, tertiary-branched C₅-C₁₈-alkyl, and

c) from 4 to 13 mol% of bivalent structural units of the formula 3



wherein a sum of molar proportion of comonomers b) and c) is between 12 and 16 mol%.

Attorney's Docket: 2002DE444
Serial No.: 10735390
Art Unit: 1714
Response to Office Action, Dated 11/15/2007

2.(Previously Presented)) The fuel oil of claim 1, wherein the molar proportion of the comonomer b) is between 5 and 11 mol%.

3.(Previously Presented) The fuel oil of claim 1, wherein the molar proportion of comonomer c) is between 4.6 and 9 mol%.

4.(Previously Presented) The fuel oil of claim 1, wherein the comonomer b) is a vinyl ester of branched carboxylic acids having from 5 to 15 carbon atoms.

5.(Previously Presented) The fuel oil of claim 1, wherein the copolymer further comprises up to 5 mol% of a further comonomer selected from the group consisting of olefins having from 3 to 18 carbon atoms, esters of acrylic acid or methacrylic acid with C₁-C₁₈-alcohols, C₁-C₁₈-alkyl vinyl ethers, and mixtures thereof.

6.(Previously Presented) The fuel oil of claim 1, wherein the copolymer has a molecular weight (by GPC against poly(styrene)) of from 3000 to 15 000 g/mol.

7.(Previously Presented) The fuel oil of claim 1, wherein the copolymer has a degree of branching determined by means of NMR between 2 and 9 CH₃/100 CH₂ groups, not taking into account the methyl groups of the comonomers.

8.(Previously Presented) The fuel oil of claim 1, wherein the copolymers have a melt viscosity at 140°C of from 20 to 10 000 mPas.

Attorney's Docket: 2002DE444
Serial No.: 10735-980
Art Unit: 1714
Response to Office Action, Dated 11/15/2007

9.(Currently Amended) The fuel oil of claim 1, wherein the total aromatic content of the mineral oil middle distillate is below 18% by weight.

10.(Currently Amended) The fuel oil of claim 1, wherein the mineral oil middle distillate has a 90-20% boiling range of less than 110°C.

11.(Currently Amended) The fuel oil of claim 1, wherein the mineral oil middle distillate has a paraffin content by DSC of more than 3% by weight of precipitated paraffins at 10°C below the cloud point.

12.(Currently Amended) The fuel oil of claim 1, wherein the mineral oil middle distillate has a density of less than 0.840 g/cm³.

13.(Currently Amended) The fuel oil of claim 1, wherein the mineral oil middle distillate additionally comprises at least one further ethylene-vinyl ester copolymer.

14.(Currently Amended) The fuel oil of claim 1, wherein the fuel oil middle distillate additionally comprises at least one polar nitrogen compound.

15.(Currently Amended) The fuel oil of claim 1, wherein the fuel oil middle distillate additionally comprises at least one alkylphenol-aldehyde resin.

Attorney's Docket: 2002DE444
 Serial No.: 10738490
 Art Unit: 1714
 Response to Office Action Dated 11/15/2007


16.(Currently Amended) The fuel oil of claim 1, wherein the fuel oil middle-distillate additionally comprises at least one comb polymer.

17.(Currently Amended) The fuel oil of claim 1, wherein the fuel oil middle-distillate additionally comprises at least one polyoxyalkylene derivative.


18.(Canceled)

19.(Currently Amended) A method for improving the cold flow behavior of a mineral oil middle distillate fuel oil, said method comprising adding a copolymer of ethylene and vinyl esters to said middle distillate having a sulfur content of at most 350 ppm, a total aromatics content of at most 22% by weight of said middle distillate, the copolymer of ethylene and vinyl esters comprising comonomers

a) bivalent structural units derived from ethylene of the formula 1




b) from 5 to 12 mol% of bivalent structural units of the formula 2



where R¹ is saturated, tertiary-branched C₅-C₁₈-alkyl, and

c) from 4 to 13 mol% of bivalent structural units of the formula 3



Attorney's Docket: 200205444
Serial No.: 10735-480
Art Unit: 1714
Response to Office Action, Dated 11/13/2007

wherein a sum of molar proportions of structural units of the formulae 2 and 3 is between 12 and 16 mol%.

20. (Previously Presented) The method of claim 19, wherein the molar proportion of the comonomer b) is between 5 and 11 mol%.

21. (Previously Presented) The method of claim 19, wherein the molar proportion of comonomer c) is between 4.6 and 9 mol%.

22. (Previously Presented) The method of claim 19, wherein the comonomer b) is a vinyl ester of branched carboxylic acids having from 5 to 15 carbon atoms.

23. (Previously Presented) The method of claim 19, wherein the copolymer further comprises up to 5 mol% of a further comonomer selected from the group consisting of olefins having from 3 to 18 carbon atoms, esters of acrylic acid or methacrylic acid with C₁-C₁₈-alcohols, C₁-C₁₈-alkyl vinyl ethers, and mixtures thereof.

24. (Previously Presented) The method of claim 19, wherein the copolymer has a molecular weight (by GPC against poly(styrene)) of from 3000 to 15 000 g/mol.

25. (Previously Presented) The method of claim 19, wherein the copolymer has a degree of branching determined by means of NMR between 2 and 9 CH₃/100 CH₂ groups, not taking into account the methyl groups of the comonomers.

Attorney's Docket: 2002DE44/
Serial No.: 10735450
Art Unit: 1714
Response to Office Action Dated 11/16/2007

26.(Previously Presented) The method of claim 19, wherein the copolymer has a melt viscosity at 140°C of from 20 to 10 000 mPas.

27.(Currently Amended) The fuel oil of claim 19, wherein the total aromatic content of the mineral oil middle distillate is below 18% by weight.


28.(Currently Amended) The method of claim 19, wherein the mineral oil middle distillate has a 90-20% boiling range of less than 110°C.

29.(Currently Amended) The method of claim 19, wherein the mineral oil middle distillate has a paraffin content by DSC of more than 3% by weight of precipitated paraffins at 10°C below the cloud point.

30.(Currently Amended) The method of claim 19, wherein the mineral oil middle distillate has a density of less than 0.840 g/cm³.


31.(New) A fuel oil comprising a proportion of mineral oil middle distillate and from 5 to 25 % by weight of a mixture of fatty acid alkyl esters, said fuel oil having a sulfur content of at most 350 ppm, a total aromatics content of at most 22% by weight of said middle distillate, and a proportion of at least one copolymer of ethylene and vinyl esters, said copolymer comprising comonomers

a) bivalent structural units derived from ethylene of the formula 1




Attorney's Docket: 2002DE444
 Serial No.: 10735-499
 Art Unit: 1714
 Response to Office Action Dated 11/15/2007

- b) from 5 to 12 mol% of bivalent structural units of the formula 2



where R¹ is a saturated, tertiary-branched C₈-C₁₈-alkyl, and

- c) from 4 to 13 mol% of bivalent structural units of the formula 3



wherein a sum of molar proportion of comonomers b) and c) is between 12 and 16 mol%.

32.(New) The fuel oil of claim 1, wherein the mixture of fatty acid alkyl esters are derived from fatty acids having from 14 to 24 carbon atoms and alcohols having from 1 to 4 carbon atoms.

33.(New) The fuel oil of claim 1, wherein the mixture of fatty acid alkyl esters is selected from the group consisting of rape seed methyl ester and mixtures of rape seed methyl esters and further vegetable oil esters.

34.(New) The fuel oil of claim 1, wherein the mixture of fatty acid methyl esters comprises rape seed methyl esters.